


Figure 1

**Add-On Treatment with Pipamperon 8-12 (mean 9) mg (bid) after Treatment with Citalopram 10-20 (mean 30) mg (bid) during 20-60 (mean 33) days (PIPCIT ADD-ON) with HDRS-17 Totalscore = 29 at Baseline in MDD in comparison with the Standard Efficacy of Antidepressants in Clinical Trials\***

**HDRS-17 REDUCTION OVER 8 WEEKS IN MDD**



A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY/VOL 57/APR 2000)

BEST AVAILABLE COPY

# HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON AS ADD-ON - CITALOPRAM vs SNRI (duloxetine) in MAJOR DEPRESSION




Figure 2

# REMISSION RATES (HDRS-17 $\leq 7$ ): COMBO PIPAMPERON AS ADD- ON - CITALOPRAM vs SNRI (venlafaxine) vs SSRI's vs PLACEBO in MAJOR DEPRESSION





Figure 3

**Foregoing Treatment During 1-5 (mean 4) days with Pipamperon 8-12 (mean 10) g/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-50 (mean 26) mg/day (bid) (PICIT FG 1-5) in MDD (HDRS-17 at BL = 23) in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials\***

### HDRS-17 REDUCTION OVER 8 WEEKS IN MDD




A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY/VOL 57/APR 2000

BEST AVAILABLE COPY

Figure 4

# HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON- CITALOPRAM WITH A FORE-GOING TREATMENT OF 4 DAYS WITH PIPAMPERON 10mg/day vs SNRI (duloxetine) in MAJOR DEPRESSION



BEST AVAILABLE COPY

Figure 5



# REMISSION RATES (HDRS-17 $\leq 7$ ): COMBO PIPAMPERON-CITALOPRAM WITH A FORE-GOING TREATMENT OF 4 DAYS WITH PIPAMPERON 10mg/day vs SNRI (venlafaxine) in MAJOR DEPRESSION





Figure 6

BEST AVAILABLE COPY

**Foregoing Treatment During 6-8 (mean 7) days with Pipamperon 8-12 (mean 11) mg/day (bid) Followed With the Combination Treatment of Pipamperon and Citalopram 20-40 (mean 30) mg/day (bid) (PPCIT FG 6-8) in MDD (HDRS-17 at BL = 28 in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials\***

**HDRS-17 REDUCTION OVER 8 WEEKS IN MDD**



A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY, VOL 57, APR 2000

**BEST AVAILABLE COPY**

Figure 7

# HDRS-17 CHANGE FROM BASELINE: COMBO PIPAMPERON- CITALOPRAM WITH A FORE-GOING TREATMENT OF 7 DAYS WITH PIPAMPERON 11mg/day vs SNRI (duloxetine) in MAJOR DEPRESSION





Figure 8



**Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in MDD in Comparison with the Standard Efficacy of Antidepressants in Clinical Trials\***

**HDRS-17 REDUCTION OVER 8 WEEKS IN MDD**



\* A. KHAN et al, Symptom Reduction and Suicide Risk in Patients Treated With Placebo in Antidepressant Clinical Trials, ARCH. OF GENERAL PSYCHIATRY / VOL 57, APR 2000)

**BEST AVAILABLE COPY**

Figure 9

# **HDRS-17 CHANGE FROM BASELINE: Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SNRI duloxetine in MAJOR DEPRESSION**




Figure 10

# REMISSION RATES (HDRS-17 $\leq 7$ ): Foregoing & Add-On Treatment with Pipamperon 8-12 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SNRI venlafaxine in MAJOR DEPRESSION





Figure 11

BEST AVAILABLE COPY

# **Y-BOCS TOTAL SCORE: Foregoing & Add-On Treatment with Pipamperon 8-16 mg/day (bid) and Citalopram 30-80 mg/day (bid) in comparison with the SSRI fluvoxamine in OCD**



COMBO PIPAMPERON-CITALOPRAM (n=7)


FLUVOXAMINE Controlled Release mean  
271 mg/day (n=253) (Hollander E, Journal of  
Clinical Psychiatry 64: 640-647, Jun 2003)

BEST AVAILABLE COPY

Figure 12



**Y-BOCS OBSESSION SCORE: Foregoing & Add-On Treatment  
with Pipamperon 8-15 mg/day (bid) and Citalopram 30-80 mg/day  
(bid) in comparison with the SSRI fluvoxamine in OCD**



—○— COMBO PIPAMPERON-CITALOPRAM (n=7)

---○--- FLUVOXAMINE Controlled Release mean  
271 mg/day (n=253) (Hollander E, Journal of  
Clinical Psychiatry 64: 640-647, Jun 2003)

Figure 13

**Y-BOCS COMPULSION SCORE: Foregoing & Add-On Treatment  
with Pipamperon 8-15 mg/day (bid) and Citalopram 30-80 mg/day  
(bid) in comparison with the SSRI fluvoxamine in OCD**





Figure 14

# **CGI-SEVERITY SCORE: Foregoing & Add-On Treatment with Pipamperon 8 mg/day (bid) and Citalopram 20-40 mg/day (bid) in comparison with the SSRI in Panic Disorder**



—○— Paroxetine in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004

—\*— COMBO PIPAMPERON-CITALOPRAM (n=3)

---○--- Sertraline in the treatment of panic disorder, Journal of Clinical Psychiatry 65: 405-413, No. 3, Mar 2004

Figure 15